

DALVANCE® Referral Form

Fax completed form to: 833-908-1122



PATIENT INFORMATION			
Patient Name:	Date of Birth:	Referral Date:	
Address:		City/State/Zip:	
Home Phone:	Cell Phone:	Work Phone:	
Secondary Contact:	Height:	Weight:	Male Female
Patient Diagnosis & ICD-10:			
Allergies:			

PROVIDER INFORMATION		
Physician Name:	Lic.#:	DEA #:
Practice Name:		NPI#:
Address:		City/State/Zip:
Office Contact:	Phone:	Fax:
Supervisory Physician (if applicable):		

PLEASE ATTACH	
Patient demographics & front/back copy of all insurance cards (prescription & medical) Recent office visit notes, history & physical, lab & pertinent procedure results Current medication list & list of prior medications tried and failed (with dates) Line access documentation/verification if applicable	Estimated creatinine clearance Culture & sensitivity results Letter of medical necessity if drug dosing or indication is outside of FDA guidelines

NURSING & LAB ORDERS
Nurse Orders: Nurse to provide assessment, teaching, lab draws, medication administration and vascular access device insertion and/or management per physician orders. Flush Orders: NaCl 0.9% - 5-10mL flush pre and post infusion and as needed Heparin - 10units/mL ---OR--- 100units/mL - 3-5mL flush after post-infusion NS flush if indicated to maintain line Lab Orders: Lab Date & Frequency:

PRESCRIPTION ORDERS			
Anaphylaxis Kit:	Epinephrine 0.3mg IM as needed	Solu-Cortef 250mg-500mg IV infusion as needed	Solu-Medrol 60mg - 125mg IV infusion as needed
(Check all that apply)	Diphenhydramine _____ mg IV infusion as needed	NS Hydration 500 ml IV infusion over 30 minutes as needed	Other

Supply Orders: All supplies for vascular access line care, drug administration kit(s), pump, and IV pole will be provided as necessary

PRODUCT	PRESCRIPTION INFORMATION	REFILLS
Is this a first dose? Yes No If No, when was last dose given? _____ When is patient due for next dose? _____		
DALVANCE <i>(to be mixed in DSW)</i>	Adult Dosing: Estimated Creatinine Clearance 30mL/min and above or on regular hemodialysis: 1500mg single dose regimen or 1000mg followed by one week later 500mg two dose regimen IV infusion via gravity ---OR--- pump over 30 minutes Less than 30mL/min and not on regular hemodialysis: 1125mg single dose regimen or 750mg followed by one week later 375mg two dose regimen IV infusion via gravity ---OR--- pump over 30 minutes	_____
OTHER		_____

By signing this form and utilizing our services, you are authorizing Amerita, Inc. to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber's Signature Dispense as Written	Print Name	Date	Prescriber's Signature Substitution Permitted	Print Name	Date
--	------------	------	---	------------	------

